

Relativistic Quark Model Calculation of the l_1, l_2 Coefficients of the Chiral Lagrangian

Felipe J. Llanes-Estrada *


Depto. Física Teórica I, Univ. Complutense, Madrid, Spain

Pedro Bicudo

Dep .Física and CFIF, Instituto Superior Técnico, 1049-001 Lisboa, Portugal

We briefly report on a relativistic quark model scheme to calculate the $O(P^4)$ pion-pion vertex in the planar approximation and in the chiral limit. The calculation is reduced to the solution of simple integral equations (Bethe-Salpeter like) by an effective use of chiral Ward Identities. Specific model computations are provided.

The interest of meson scattering in the quark model has recently revived due to the exotic 1^{-+} mesons seen in BNL at the energies of 1.4 and 1.6 *GeV*. The hybrid [1] or meson molecule nature of these states is being debated. The standard tools for meson scattering are the chiral lagrangian [2] and its unitarized extensions [3], which cannot access that intermediate energy region, let alone interpret the Fock space nature of the resonances. Still they provide the benchmark for model approaches to the underlying dynamics which encompass chiral symmetry. In recent works [4, 5] the use of a novel chiral Ward Identity (3 below) allowed the reduction of the planar quark model approximation to pion scattering,




to reproduce the $O(P^2)$, $O(m_\pi^2)$ chiral amplitude. Here we continue this effort by establishing contact with the $O(p^4)$, $O(m_\pi^0)$ chiral lagrangian. This work is related to microscopic studies of $\pi - \pi$ scattering with the Nambu and Jona-Lasinio model [7], with the bosonization technique [8], with the equal time quark model [4, 9], and with a direct numerical evaluation with a Schwinger-Dyson model [4, 10]. While the $O(P^2)$ amplitude is model independent, the $O(p^4)$ provides a model test. Here we exactly and explicitly determine the Feynman diagrams that govern the $O(p^4)$ amplitude, then employ two simple models for a test evaluation.

In the standard notation, the pion scattering amplitude arising from its $O(P^4)$ contact terms alone reads

$$A(s, t, u)^{(P^4)} = \frac{1}{F^4} \left[\left(2l_1 + \frac{l_2}{2} \right) s^2 + \frac{l_2}{2} (t - u)^2 \right]. \quad (2)$$

The Bethe-Salpeter pion vertex χ_π in (1) above, function of two momenta (incoming meson P and quark $k + P/2$) can be conveniently expanded in an external momentum power series. But first it is useful to extract from it a dressed vertex Γ_A satisfying interesting Ward Identities connecting it to the fermion dressed propagator and the planar ladder,

$$\chi(P, k) = \frac{-i\Gamma_A(P, k) + \Delta(P, k)}{\sqrt{2}f_\pi} \quad (3)$$

$$\Gamma_A(P, k + P/2) = S^{-1}(k - P/2)\gamma_5 + \gamma_5 S^{-1}(k + P/2)$$


* Talk presented by F. Llanes-Estrada. Thanks to E. Ribeiro, P. Maris, S. Cotanch, A. Szczepaniak and J. Pelaez for valuable input. This work supported by an Univ. Complutense travel allowance, and project FPA 2000-0956.

The use of these identities, as well as the fact that no zero-momentum pole appears in any vector or scalar ladder given the symmetry breaking pattern of the theory, allows us to reduce the momentum four term of eq. (1) to three different classes of diagrams, parameterized as,

$$\begin{aligned}
 & \begin{array}{c} \Delta_1 \quad \Delta_2 \\ \text{---} \text{---} \text{---} \text{---} \\ \Delta_4 \quad \Delta_3 \end{array} + \begin{array}{c} \Delta_4 \quad \Delta_1 \\ \text{---} \text{---} \text{---} \text{---} \\ \Delta_3 \quad \Delta_2 \end{array} - \begin{array}{c} \Delta_4 \quad \Delta_1 \\ \text{---} \text{---} \text{---} \text{---} \\ \Delta_3 \quad \Delta_2 \end{array} \\
 &= 3d_1(P_1 \cdot P_2 \ P_3 \cdot P_4 + P_1 \cdot P_4 \ P_2 \cdot P_3) + 3d_2 P_1 \cdot P_3 \ P_2 \cdot P_4 , \\
 & \begin{array}{c} k \\ \text{---} \text{---} \text{---} \text{---} \\ P_3 \quad \Delta_1 \\ k-P_1-P_2 \quad \Delta_2 \end{array} = 3d_3 P_1 \cdot P_3 \ P_1 \cdot P_2 + 3d_4 P_2 \cdot P_3 \ P_1 \cdot P_2 , \\
 & \begin{array}{c} q \\ \text{---} \text{---} \text{---} \text{---} \\ P_1 \quad \Delta_1 \\ q+P_1+P_4 \quad \Delta_2 \end{array} = 3d_5 P_1 \cdot P_3 \ P_1 \cdot P_4 + 3d_6 P_3 \cdot P_4 \ P_1 \cdot P_4 .
 \end{aligned} \tag{4}$$

All external momentum and color dependence is explicit. To calculate the ladders which remain in eq. (4), we write integral equations for them, analogous to the Bethe-Salpeter equation for a meson and with up to ten independent functions of q^2 . Matching this Feynman amplitude with the physical external momenta and multiplying each permutation by the adequate isospin factors, we obtain for the l 's defined in eq. (2),

$$\begin{aligned}
 l_1 &= \frac{3}{32}(-2d_1 + d_2 - 6d_3 + 2d_4 + 5d_5 + d_6) \\
 l_2 &= -\frac{3}{16}(d_2 - 2d_3 + 2d_4 + d_5 - d_6) .
 \end{aligned} \tag{5}$$

The necessary Dirac gamma traces are computed with FORM [6]. We perform model evaluations in Euclidean space with an interaction rung given by $V = -g^2 \gamma_\mu iK(k - q) \gamma^\mu$ which is similar to the One Gluon Exchange interaction in the Feynman Gauge. We have used two different infrared strong kernels, both UV finite to avoid renormalization complications.

The first is a simple Gaussian kernel (whose Euclidean angular integrals are simply Bessel functions) $K(q) = \exp(-q^2/\Lambda^2)$ with parameters $g = 3$, $\Lambda = 1.15 \text{ GeV}$ (model 1) which leads to a constituent quark mass of 116 MeV , a condensate $-(170 \text{ MeV})^3$, a pion decay constant $f_\pi = 87 \text{ MeV}$ and gives $l_1 = -0.11$, $l_2 = 0.14$; while the parameters (model 2) $g = 1.7$, $\Lambda = 2$ lead to a mass of 93 MeV , condensate $-(229 \text{ MeV})^3$, and $f_\pi = 86 \text{ MeV}$, $l_1 = -0.06$, $l_2 = 0.12$. The second is a rational kernel (whose Euclidean angular integrals contain only polynomials and square roots) $K(q) = [1/(q^2 - \lambda^2) - 1/(q^2 - \Lambda^2)]$ with $g = 10.3$, $\lambda = 0.8 \text{ GeV}$, $\Lambda = 0.85 \text{ GeV}$ (model 3), we obtain a constituent quark mass of 216 MeV , condensate $-(156 \text{ MeV})^3$, pion decay constant 92 MeV , and $l_1 = -0.065$, $l_2 = -0.12$.

The computed l_1 and l_2 have the correct signs and ratio when compared with the phenomenological analysis, however they are large. Investigations are underway to test a wider class of quark-antiquark interactions.

-
- [1] F. J. Llanes-Estrada and S. R. Cotanch, Phys. Lett. **B504**, 15 (2001).
 - [2] J. Gasser and H. Leutwyler, Ann. Phys. 158, 142 (1984).
 - [3] A. Gomez Nicola and J. R. Pelaez, Phys. Rev. **D65**:054009 (2002); A. Dobado, M. J. Herrero and T. N. Truong, Phys. Lett. **B235**, 134 (1990).
 - [4] P. Bicudo *et al.* Phys. Rev. **D65**: 076008 (2002).
 - [5] P. Bicudo, in *Quark Confinement and the Hadron Spectrum V*, World Scientific, ed. N. Brambilla and G. Prosperi (2002) [arXiv:nuc-th/0110052].
 - [6] J. A. M. Vermaseren "New Features of Form", math-ph/0010025 preprint.
 - [7] V. Bernard *et al.* Phys. Lett. **B253**, 443 (1991).
 - [8] C. Roberts *et al.* Phys. Rev. D **49**, 125 (1994).
 - [9] P. Bicudo, M. Faria, G. Marques, and J. Ribeiro [arXiv:nucl-th/0106071].
 - [10] S. R. Cotanch and P. Maris, [arXiv:hep-ph/0210151].




FIG. 1: Mass gap function solution of the Schwinger-Dyson equation for model 1 (in GeV).




FIG. 2: Solution of the homogeneous Bethe-Salpeter equation. From top to bottom, $F0[a]$, $E2[q]$, $G0[q]$ (corresponding to the coefficients of an external momentum expansion of the llewelyn-Smith wavefunctions).




FIG. 3: Solution of the homogeneous equation for two Δ 's with ladder, needed to calculate $d1, d2, d3, d4$. (Coefficients of external momentum expansion).




FIG. 4: Solution of the homogeneous equation for P with ladder, needed to calculate $d5, d6$. (Coefficients of external momentum expansion).